

# MAJUMDAR-PAPAPETROU TYPE SOLUTIONS IN SIGMA-MODEL AND INTERSECTING P-BRANES

V.D. Ivashchuk<sup>1</sup> and V.N. Melnikov<sup>2</sup>

*Center for Gravitation and Fundamental Metrology, VNIIMS  
3-1, M.Ulyanovoy Str., Moscow 117313, Russia*

## Abstract

The block-orthogonal generalization of the Majumdar-Papapetrou type solutions for the  $\sigma$ -model studied earlier are obtained and corresponding solutions with  $p$ -branes are considered. The existence of solutions and the number of independent harmonic functions is defined by the matrix of scalar products of vectors  $U^s$ , governing the  $\sigma$ -model target space metric. For orthogonal  $U^s$ , when target space is a symmetric homogeneous space, the solutions reduce to the previous ones. Two special classes of obtained solutions with  $U^s$  related to finite dimensional Lie algebras and hyperbolic (Kac-Moody) algebras are singled out and investigated. The affine Cartan matrices do not arise in the scheme under consideration. Some examples of solutions and intersection rules for  $D = 11$  supergravity, related  $D = 12$  theory and extending them  $B_D$ -models are considered. For special multicenter solutions criterions for the existence of horizon and curvature singularity are found.

---

<sup>1</sup>ivas@rgs.phys.msu.su

<sup>2</sup>rgs@com2com.ru

# 1 Introduction

In this paper we consider a family of Majumdar-Papapetrou (MP) type solutions [1] for gravitating sigma-model (for review see, for example, [2]) with the action

$$S_\sigma = \int d^{d_0}x \sqrt{|g^0|} \left\{ R[g^0] - \hat{G}_{AB} g^{0\mu\nu} \partial_\mu \sigma^A \partial_\nu \sigma^B - 2V(\sigma) - \sum_{s \in S} \varepsilon_s e^{-2U_A^s \sigma^A} g^{0\mu\nu} \partial_\mu \Phi^s \partial_\nu \Phi^s \right\}, \quad (1.1)$$

where  $g^0 = g_{\mu\nu}^0(x) dx^\mu \otimes dx^\nu$  is a metric on  $d_0$ -dimensional manifold  $M_0$ ,  $R[g^0]$  is the scalar curvature of  $g^0$ ,  $\sigma = (\sigma^A) \in \mathbf{R}^N$  and  $\Phi = (\Phi^s, s \in S)$  is a set of scalar fields ( $S \neq \emptyset$ ),  $V = V(\sigma)$  is a potential,  $(\hat{G}_{AB})$  is a symmetric non-degenerate matrix,  $U^s = (U_A^s) \in \mathbf{R}^N$  are vectors and  $\varepsilon_s = \pm 1$ ,  $s \in S$ . The sigma-model (1.1) arises in multidimensional gravitational models with scalar fields and fields of forms, when solutions with intersecting  $p$ -branes are considered [3]-[6] (a pure gravitational sector of the sigma-model was considered in [7, 8, 9]). For  $p$ -brane applications (see, for example, [3]-[35])  $g^0$  is Euclidean,  $(\hat{G}_{AB})$  is positively defined and  $\varepsilon_s = -1$ , if pseudo-Euclidean (electric and magnetic)  $p$ -branes in a pseudo-Euclidean space-time are considered. The sigma-model (1.1) may be also considered for the pseudo-Euclidean metric  $g^0$  of signature  $(-, +, \dots, +)$ . In this case for a positively defined matrix  $(\hat{G}_{AB})$  and  $\varepsilon_s = 1$  we get a non-negative kinetic energy terms.

The target space of the model is  $(\mathbf{R}^K, \mathcal{G})$ , where

$$\mathcal{G} = \hat{G}_{AB} d\sigma^A \otimes d\sigma^B + \sum_{s \in S} \varepsilon_s e^{-2U_A^s \sigma^A} d\Phi^s \otimes d\Phi^s, \quad (1.2)$$

It was proved in [38] that the target space  $\mathcal{T} = (\mathbf{R}^K, \mathcal{G})$  is a homogeneous (coset) space  $G/H$  ( $G$  is the isometry group of  $\mathcal{T}$ ,  $H$  is the isotropy subgroup of  $G$ ).  $\mathcal{T}$  is symmetric (i.e. the Riemann tensor is covariantly constant:  $\nabla_M R_{M_1 M_2 M_3 M_4}[\mathcal{G}] = 0$ ) if and only if

$$(U^{s_1} - U^{s_2})(U^{s_1}, U^{s_2}) = 0 \quad (1.3)$$

for all  $s_1, s_2 \in S$ , i.e. when any two vectors  $U^{s_1}$  and  $U^{s_2}$ ,  $s_1 \neq s_2$ , are either coinciding  $U^{s_1} = U^{s_2}$  or orthogonal  $(U^{s_1}, U^{s_2}) = 0$ , where scalar product  $(\cdot, \cdot)$  is defined as follows

$$(U, U') = \hat{G}^{AB} U_A U'_B, \quad (1.4)$$

for  $U, U' \in \mathbf{R}^N$ , where  $(\hat{G}^{AB}) = (\hat{G}_{AB})^{-1}$ .

In our previous papers [3]-[6] we considered the orthogonal case

$$(U^{s_1}, U^{s_2}) = 0, \quad (1.5)$$

$s_1 \neq s_2$ , and  $(U^{s_1}, U^{s_1}) \neq 0$ ,  $s_1, s_2 \in S$ , and for the zero-potential:  $V = 0$ , obtained a family of exact solutions in the sigma-model (1.1) governed by  $k$  harmonic functions, where  $k$  is a number of  $s$ , satisfying  $\varepsilon_s(U^s, U^s) < 0$ . Using these sigma-model solutions we also found a family of solutions in multidimensional gravity with  $p$ -branes. For  $p$ -brane applications the orthogonality condition (1.5) is equivalent to (orthogonal) intersection rules [3, 4, 5, 6, 19, 20] also known as no-force condition.

In [5] these solutions (sigma-model and  $p$ -brane ones) were also generalized to the case  $V \neq 0$ , where  $V = \sum_{a=1}^m A_a e^{u_a^a \sigma^A}$  and  $(u^a, U^s) = 0$ ,  $a = 1, \dots, m$ ,  $s \in S$ .

Here we generalize the orthogonal  $\sigma$ -model solutions from [3]-[6] to a more general block-orthogonal case:

$$S = S_1 \cup \dots \cup S_k, \quad S_i \cap S_j = \emptyset, \quad i \neq j \quad (1.6)$$

$S_i \neq \emptyset$ , i.e. the set  $S$  is a union of  $k$  non-intersecting (non-empty) subsets  $S_1, \dots, S_k$ , and

$$(U^s, U^{s'}) = 0 \quad (1.7)$$

for all  $s \in S_i$ ,  $s' \in S_j$ ,  $i \neq j$ ;  $i, j = 1, \dots, k$ . According to (1.7) the set of vectors  $(U^s, s \in S)$  has a block-orthogonal structure with respect to the scalar product (1.4): it is splitted into  $k$  mutually orthogonal blocks  $(U^s, s \in S_i)$ ,  $i = 1, \dots, k$ .

In this case solutions do exist, when the matrix of scalar products  $B = ((U^{s_1}, U^{s_2}))$  and  $\varepsilon_s$  satisfy the relation

$$\sum_{s' \in S} (U^s, U^{s'}) \varepsilon_{s'} \nu_{s'}^2 = -1 \quad (1.8)$$

for some set of real  $\nu_s$ ,  $s \in S$ . The number of independent harmonic functions is defined by the number of blocks in the matrix  $B$ .

The paper is organized as follows. In Sect. 2 the sigma-model is considered and in block-orthogonal case exact solutions governed by a set of harmonic functions are obtained. In Sect. 3 certain examples of solutions related to Lie algebras (e.g. finite dimensional and hyperbolic) are considered and restrictions on signature parameters  $\varepsilon_s$  for different Lie algebras are derived. It is shown that affine Kac-Moody algebras do not appear in this scheme. Sect. 4 is devoted to application of the sigma-model solutions to multidimensional models with intersecting p-branes. The block-orthogonal generalization of (orthogonal) p-brane MP type solutions is presented in subsect. 4.2 (We note, that recently block-orthogonal black hole and wormhole solutions with p-branes were considered in [33]. These solutions generalize orthogonal black hole solutions [24, 25, 26, 27, 32, 35].) In subsect. 4.3 the behavior of the Riemann tensor squared (Kretschmann scalar) for multicenter solutions is investigated and criteria for the existence of horizon and finiteness of the Kretschmann scalar are established. In Sect. 5 intersection rules and some examples are considered (e.g. for  $B_D$ -models that may be relevant for future generalizations of  $M$ - and  $F$ -theories [28, 29] to dimensions  $D > 12$ .) Here a dyon solution for  $D = 11$  supergravity with electric 2-brane and magnetic 5-brane configuration is considered. This solution has  $A_2 = sl(3, \mathbf{C})$  intersection rule:  $3 \cap 6 = 1$  instead of the orthogonal one  $3 \cap 6 = 2$ . An analogous dyon solution for  $D = 10$  IIA supergravity was recently considered in [37].

We note that the intersecting  $p$ -brane solutions (see [3, 4, 5, 6, 17, 19, 20, 21] and references therein) with “orthogonal” intersection rules correspond to the Lie algebras  $A_1 \oplus \dots \oplus A_1$ , where  $A_1 = sl(2, \mathbf{C})$ . (The MP solution in this classification corresponds to the algebra  $A_1$ .) In supergravitational models these solutions correspond to the so-called BPS saturated states preserving fractional supersymmetry [11]-[22].

## 2 $\sigma$ -model solutions

Here we consider a family of solutions to equations of motion of the sigma-model (1.1) in the block-orthogonal case (1.6), (1.7).

Equations of motion corresponding to (1.1) have the following form

$$R_{\mu\nu}[g^0] = \hat{G}_{AB} \partial_\mu \sigma^A \partial_\nu \sigma^B + \frac{2V}{d_0 - 2} g_{\mu\nu}^0 + \sum_{s \in S} \varepsilon_s e^{-2U_A^s \sigma^A} \partial_\mu \Phi^s \partial_\nu \Phi^s, \quad (2.1)$$

$$\hat{G}_{AB} \Delta[g^0] \sigma^B - \frac{\partial V}{\partial \sigma^A} + \sum_{s \in S} \varepsilon_s U_A^s e^{-2U_C^s \sigma^C} g^{0\mu\nu} \partial_\mu \Phi^s \partial_\nu \Phi^s = 0, \quad (2.2)$$

$$\partial_\mu \left( \sqrt{|g^0|} g^{0\mu\nu} e^{-2U_A^s \sigma^A} \partial_\nu \Phi^s \right) = 0, \quad (2.3)$$

$s \in S$ . Here  $\Delta[g^0]$  is the Laplace-Beltrami operator corresponding to  $g^0$ .

**Proposition 1.** Let  $(M_0, g^0)$  be Ricci-flat

$$R_{\mu\nu}[g^0] = 0. \quad (2.4)$$

Then the field configuration

$$g^0, \quad \sigma^A = \sum_{s \in S} \varepsilon_s U^{sA} \nu_s^2 \ln H_s, \quad \Phi^s = \frac{\nu_s}{H_s}, \quad (2.5)$$

$s \in S$ , satisfies the field equations (2.1)–(2.3) with  $V = 0$  if real numbers  $\nu_s$  obey the relations

$$\sum_{s' \in S} (U^s, U^{s'}) \varepsilon_{s'} \nu_{s'}^2 = -1$$

$s \in S$ , functions  $H_s > 0$  are harmonic, i.e.

$$\Delta[g^0]H_s = 0, \quad (2.6)$$

$s \in S$ , and  $H_s$  are coinciding inside blocks:

$$H_s = H_{s'} \quad (2.7)$$

for  $s, s' \in S_i$ ,  $i = 1, \dots, k$ .

The Proposition 1 can be readily verified by a straightforward substitution of (2.4)–(2.7) into equations of motion (2.1)–(2.3). In special (orthogonal) case, when any block contains only one vector (i.e. all  $|S_i| = 1$ ) the Proposition 1 coincides with that of [5]. In general case vectors inside each block  $S_i$  are not orthogonal. The solution under consideration depends on  $k$  independent harmonic functions. For a given set of vectors  $(U^s, s \in S)$  the maximal number  $k$  arises for irreducible block-orthogonal decomposition (1.6), (1.7), when any block  $(U^s, s \in S_i)$  can not be splitted into two mutually-orthogonal subblocks.

We note that due to (1.7) the relation (1.8) may be rewritten as

$$\sum_{s' \in S_i} (U^s, U^{s'}) \varepsilon_{s'} \nu_{s'}^2 = -1, \quad (2.8)$$

$s \in S_i$ ,  $i = 1, \dots, k$ . Hence, parameters  $(\nu_s, s \in S_i)$  depend upon vectors  $(U^s, s \in S_i)$ ,  $i = 1, \dots, k$ .

### 3 Parameters of solutions related to Lie algebras

Here we put

$$(U^s, U^s) \neq 0 \quad (3.1)$$

for all  $s \in S$  and introduce quasi-Cartan matrix  $A = (A^{ss'})$

$$A^{ss'} \equiv \frac{2(U^s, U^{s'})}{(U^{s'}, U^{s'})}, \quad (3.2)$$

$s, s' \in S$ , which coincides with the Cartan matrix, when  $U^s$ ,  $s \in S$ , are simple roots of some Lie algebra and  $(.,.)$  is standard bilinear form on the root space. From (1.7) we get a block-orthogonal structure of  $A$ :

$$A = \begin{pmatrix} A_{(1)} & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & A_{(k)} \end{pmatrix}, \quad (3.3)$$

where  $A_{(i)} = (A^{ss'}, s, s' \in S_i)$ ,  $i = 1, \dots, k$ . Here we tacitly assume that the set  $S$  is ordered,  $S_1 < \dots < S_k$  and the order in  $S_i$  is inherited by the order in  $S$ .

For  $\det A_{(i)} \neq 0$  relation (2.8) may be rewritten in the equivalent form

$$\varepsilon_s \nu_s^2 (U^s, U^s) = -2 \sum_{s' \in S_i} A_{ss'}^{(i)}, \quad (3.4)$$

$s \in S_i$ , where  $(A_{ss'}^{(i)}) = A_{(i)}^{-1}$ . Thus, eq. (2.8) may be resolved in terms of  $\nu_s$  for certain  $\varepsilon_s = \pm 1$ ,  $s \in S_i$ . For  $\det A_{(i)} = 0$  there exist situations when eq. (2.8) has no solutions even for complex  $\nu_s$ . Indeed, let us suppose that there exists a vector  $a = (a_s, s \in S_i)$  satisfying the relations

$$\sum_{s \in S_i} a_s A_{(i)}^{ss'} = 0, \quad \sum_{s \in S_i} a_s \neq 0, \quad (3.5)$$

$s' \in S_i$  (eqs. (3.5) imply  $\det A_{(i)} = 0$  and, hence,  $\det A = 0$ ). From (2.8) and the first relation in (3.5) we get  $\sum_{s \in S_i} a_s = 0$ , that contradicts the second relation in (3.5).

In what follows we consider the block-orthogonal decomposition to be irreducible, i.e. for any  $i$  the block  $(U^s, s \in S_i)$  can not be splitted into two mutually orthogonal subblocks. In this case any matrix  $A_{(i)}$  is indecomposable (or irreducible) in the sense that there is no renumbering of vectors which would bring  $A_{(i)}$  to the block diagonal form  $A_i = \text{diag}(A'_{(i)}, A''_{(i)})$ .

Let  $A$  be a generalized Cartan matrix [39, 40]. In this case

$$A^{ss'} \in -\mathbf{Z}_+ \equiv \{0, -1, -2, \dots\} \quad (3.6)$$

for  $s \neq s'$  and  $A$  generates generalized symmetrizable Kac-Moody algebra [39, 40].

Now we fix  $i \in \{1, \dots, k\}$ . From (3.3) and (3.6) we get

$$A_{(i)}^{ss'} \in -\mathbf{Z}_+, \quad (3.7)$$

$s, s' \in S_i$ ,  $s \neq s'$ . There are three possibilities for  $A_{(i)}$ : when a)  $\det A_{(i)} > 0$ , b)  $\det A_{(i)} < 0$  and c)  $\det A_{(i)} = 0$ . For  $\det A_{(i)} \neq 0$  the corresponding Kac-Moody algebra is simple, since  $A_{(i)}$  is indecomposable [40]. Now we analyze these three possibilities.

### 3.1 Finite dimensional Lie algebras

Let  $\det A_{(i)} > 0$ . In this case  $A_{(i)}$  is a Cartan matrix of a simple finite-dimensional Lie algebra and  $A_{(i)}^{ss'} \in \{0, -1, -2, -3\}$ ,  $s \neq s'$ . The elements of inverse matrix  $A_{(i)}^{-1}$  are positive (see Ch.7 in [40]) and hence we get from (3.4)

$$\varepsilon_s (U^s, U^s) < 0, \quad (3.8)$$

$s \in S_i$ .

**Example 1.** Let us consider the Cartan matrix

$$A_{(i)} = \begin{pmatrix} 2 & -q \\ -1 & 2 \end{pmatrix}, \quad (3.9)$$

$q = 1, 2, 3$ , corresponding to the Lie algebras  $A_2 = \mathfrak{sl}(3)$ ,  $B_2 = \mathfrak{so}(5)$  and  $G_2$  respectively. Relations (3.4) read in this case

$$\varepsilon_s \nu_s^2(U^s, U^s)(q - 4) = 4 + 2q, \quad 6 \quad (3.10)$$

for  $s = 1, 2$ . Here  $S_i = \{1, 2\}$ .

**Example 2.** Let  $A_{(i)}$  be  $r \times r$  Cartan matrix for the Lie algebra  $A_r = \mathfrak{sl}(r + 1)$ ,  $r \geq 2$ . This matrix is described graphically by the Dynkin diagram pictured on Fig.1.




Fig.1. *Dynkin diagram for  $A_r$  Lie algebra*

(For  $s \neq s'$ ,  $A_{(i)}^{ss'} = -1$  if nodes  $s$  and  $s'$  are connected by a line on the diagram and  $A_{(i)}^{ss'} = 0$  otherwise). Using the relation for the inverse matrix  $A_{(i)}^{-1} = (A_{ss'}^{(i)})$  (see Ch.7 in [40])

$$A_{ss'}^{(i)} = \frac{1}{r+1} \min(s, s') [r+1 - \max(s, s')] \quad (3.11)$$

we may rewrite (3.4) as follows

$$\varepsilon_s \nu_s^2(U^s, U^s) = s(s-1-r), \quad (3.12)$$

$$s \in \{1, \dots, r\} = S_i.$$

### 3.2 Hyperbolic Kac-Moody algebras

Now we consider the case  $\det A_{(i)} < 0$ . Among such irreducible symmetrizable matrices satisfying (3.7) there exists a large subclass of Cartan matrices, corresponding to infinite-dimensional simple hyperbolic generalized Kac-Moody (KM) algebras of ranks  $r = 2, \dots, 10$  [39, 40].

**Example 3.** Let

$$A_{(i)} = \begin{pmatrix} 2 & -q_1 \\ -q_2 & 2 \end{pmatrix}, \quad q_1 q_2 > 4, \quad (3.13)$$

$q_1, q_2 \in \mathbf{N}$ . This is the Cartan matrix for the hyperbolic KM algebra  $H_2(q_1, q_2)$ . From (3.4) we get

$$\varepsilon_s \nu_s^2(U^s, U^s)(q_1 q_2 - 4) = 2q_s + 4, \quad (3.14)$$

$$s \in \{1, 2\} = S_i.$$

**Example 4.** Let  $A_{(i)}$  be a Cartan matrix corresponding to  $E_{10}$  hyperbolic KM algebra with the Dynkin diagram pictured on Fig.2.




Fig.2. *Dynkin diagram for  $E_{10}$  hyperbolic KM algebra*

In this case we get from (3.4) [37]

$$\frac{1}{2}\varepsilon_s(U^s, U^s)\nu_s^2 = 30, 61, 93, 126, 160, 195, 231, 153, 76, 115 \quad (3.15)$$

for  $s = 1, 2, \dots, 10$  respectively.

In both examples for hyperbolic algebras the following relations are satisfied

$$\varepsilon_s(U^s, U^s) > 0, \quad (3.16)$$

$s \in S_i$ . This relation is valid in the general case, since  $(A_{(i)}^{-1})_{ss'} \leq 0$ ,  $s, s' \in S$ , for any hyperbolic algebra [41].

Hyperbolic KM algebras appeared in different areas of mathematical physics, e.g. in ordinary gravity [42] ( $\mathcal{F}_3$  hyperbolic algebra), supergravity: [43, 44] ( $E_{10}$  hyperbolic algebra), [45] ( $\mathcal{F}_3$  hyperbolic algebra), strings etc (see also [46] and references therein). In [45] it was shown that the chiral reduction of a simple ( $N = 1$ ) supergravity from four dimensions to one dimension gives rise to the hyperbolic algebra of rank 3 (namely  $\mathcal{F}_3$ ). In [36] an example of cosmological solution in  $D = 11$  supergravity describing three Euclidean  $p$ -branes (two magnetic and one electric) with intersection rules corresponding to the hyperbolic KM algebra  $\mathcal{F}_3$  was constructed.

### 3.3 Affine Kac-Moody algebras

Now we proceed to the degenerate case:  $\det A_{(i)} = 0$ . Here we restrict ourselves to a subclass of affine KM algebras [39, 40]). Unfortunately, there are no solutions considered above in the affine case. Indeed, any affine Cartan matrix satisfy the relations (3.5) with  $a_s > 0$  called Coxeter labels and, hence, the solutions are absent in this case.

## 4 Solutions with intersecting $p$ -branes

### 4.1 The model

Now we consider a multidimensional gravitational model governed by the action [5]

$$S = \int d^D z \sqrt{|g|} \left\{ R[g] - h_{\alpha\beta} g^{MN} \partial_M \varphi^\alpha \partial_N \varphi^\beta - \sum_{a \in \Delta} \frac{\theta_a}{n_a!} \exp[2\lambda_a(\varphi)] (F^a)^2 \right\} \quad (4.1)$$

where  $g = g_{MN} dz^M \otimes dz^N$  is the metric,  $\varphi = (\varphi^\alpha) \in \mathbf{R}^l$  is a vector of scalar fields,  $(h_{\alpha\beta})$  is a symmetric non-degenerate  $l \times l$  matrix ( $l \in \mathbf{N}$ ),  $\theta_a = \pm 1$ ,  $F^a = dA^a$  is  $n_a$ -form ( $n_a \geq 1$ ),  $\lambda_a$  is a 1-form on  $\mathbf{R}^l$ :  $\lambda_a(\varphi) = \lambda_{\alpha a} \varphi^\alpha$ ,  $a \in \Delta$ ,  $\alpha = 1, \dots, l$ . Here  $\Delta$  is some finite set. In the models with one time all  $\theta_a = 1$  when the signature of the metric is  $(-1, +1, \dots, +1)$ .

We consider the manifold

$$M = M_0 \times M_1 \times \dots \times M_n, \quad (4.2)$$

with the metric

$$g = e^{2\gamma(x)} g^0 + \sum_{i=1}^n e^{2\phi^i(x)} g^i \quad (4.3)$$

where  $g^0 = g_{\mu\nu}^0(x)dx^\mu \otimes dx^\nu$  is a metric on the manifold  $M_0$ , and  $g^i = g_{m_i n_i}^i(y_i)dy_i^{m_i} \otimes dy_i^{n_i}$  is a metric on the manifold  $M_i$  satisfying

$$Ric[g^i] = \xi_i g^i, \quad (4.4)$$

$\xi_i = \text{const}$ ,  $i = 1, \dots, n$  ( $Ric[g]$  denotes Ricci-tensor, corresponding to  $g$ ). Thus, all internal spaces  $(M_i, g^i)$ ,  $i = 1, \dots, n$ , are Einstein ones. Any manifold  $M_\nu$  is claimed to be oriented and connected and  $d_\nu \equiv \dim M_\nu$ ,  $\nu = 0, \dots, n$ . Let

$$\tau_i \equiv \sqrt{|g^i(y_i)|} dy_i^1 \wedge \dots \wedge dy_i^{d_i}, \quad \varepsilon(i) \equiv \text{sign}(\det(g_{m_i n_i}^i)) = \pm 1 \quad (4.5)$$

denote the volume  $d_i$ -form and signature parameter respectively,  $i = 1, \dots, n$ . Let  $\Omega = \Omega_n$  be a set of all subsets of  $\{1, \dots, n\}$ ,  $|\Omega| = 2^n$ . For any  $I = \{i_1, \dots, i_k\} \in \Omega$ ,  $i_1 < \dots < i_k$ , we denote

$$\tau(I) \equiv \tau_{i_1} \wedge \dots \wedge \tau_{i_k}, \quad d(I) \equiv \sum_{i \in I} d_i, \quad \varepsilon(I) \equiv \prod_{i \in I} \varepsilon(i). \quad (4.6)$$

We also put  $\tau(\emptyset) = \varepsilon(\emptyset) = 1$  and  $d(\emptyset) = 0$ .

For fields of forms we consider the following composite electromagnetic ansatz

$$F^a = \sum_{I \in \Omega_{a,e}} \mathcal{F}^{(a,e,I)} + \sum_{J \in \Omega_{a,m}} \mathcal{F}^{(a,m,J)} \quad (4.7)$$

where

$$\mathcal{F}^{(a,e,I)} = d\Phi^{(a,e,I)} \wedge \tau(I), \quad (4.8)$$

$$\mathcal{F}^{(a,m,J)} = e^{-2\lambda_a(\varphi)} * (d\Phi^{(a,m,J)} \wedge \tau(J)) \quad (4.9)$$

are elementary forms of electric and magnetic types respectively,  $a \in \Delta$ ,  $I \in \Omega_{a,e}$ ,  $J \in \Omega_{a,m}$  and  $\Omega_{a,e} \subset \Omega$ ,  $\Omega_{a,m} \subset \Omega$ . In (4.9)  $*$  =  $*[g]$  is the Hodge operator on  $(M, g)$ . For scalar functions we put

$$\varphi^\alpha = \varphi^\alpha(x), \quad \Phi^s = \Phi^s(x), \quad (4.10)$$

$s \in S$ . Here and below

$$S = S_e \cup S_m, \quad S_v = \bigcup_{a \in \Delta} \{a\} \times \{v\} \times \Omega_{a,v}, \quad (4.11)$$

$v = e, m$ .

Due to (4.8) and (4.9)

$$d(I) = n_a - 1, \quad d(J) = D - n_a - 1, \quad (4.12)$$

for  $I \in \Omega_{a,e}$ ,  $J \in \Omega_{a,m}$ .

**Remark 1.** It is more correct to write in (4.3)  $\hat{g}^i$  instead of  $g^i$ , where  $\hat{g}^i = p_i^* g^i$  is the pullback of the metric  $g^i$  to the manifold  $M$  by the canonical projection:  $p_i : M \rightarrow M_i$ ,  $i = 1, \dots, n$ . Here we omit all “hats” (e.g. corresponding to volume  $\tau$ -forms) for simplicity.

Let  $d_0 \neq 2$  and

$$\gamma = \gamma_0(\phi) \equiv \frac{1}{2 - d_0} \sum_{j=1}^n d_j \phi^j, \quad (4.13)$$

i.e. the generalized harmonic gauge is used.

Now we impose restriction on sets  $\Omega_{a,v}$ . These restrictions guarantee the block-diagonal structure of a stress-energy tensor (like for the metric) and the existence of  $\sigma$ -model representation [5].

We denote  $w_1 \equiv \{i | i \in \{1, \dots, n\}, d_i = 1\}$ , and  $n_1 = |w_1|$  (i.e.  $n_1$  is the number of 1-dimensional spaces among  $M_i$ ,  $i = 1, \dots, n$ ). We also denote by  $I \sqcup J$  the union of non-intersecting sets  $I$  and  $J$

**Restriction 1.** For any  $a \in \Delta$ ,  $v \in \{e, m\}$ , there are no  $I, J \in \Omega_{a,v}$  such that

$$I = \{i\} \sqcup (I \cap J), \quad J = (I \cap J) \sqcup \{j\} \quad (4.14)$$

for some  $i, j \in w_1$ ,  $i \neq j$ .

**Restriction 2** (only for  $d_0 = 1, 3$ ). For any  $a \in \Delta$  there are no  $I \in \Omega_{a,m}$ ,  $J \in \Omega_{a,e}$  such that

$$\bar{I} = \{i\} \sqcup J \quad (4.15)$$

for  $d_0 = 1$  and

$$J = \{i\} \sqcup \bar{I} \quad (4.16)$$

for  $d_0 = 3$ , where  $i \in w_1$  and  $\bar{I}$  is defined as follows

$$\bar{I} \equiv \{1, \dots, n\} \setminus I. \quad (4.17)$$

Restriction 1 is satisfied for  $n_1 \leq 1$  and in the case when  $|\Omega_{a,v}| \leq 1$  for all  $a \in \Delta$ ,  $v \in \{e, m\}$  (e.g. in the non-composite case). For  $n_1 \geq 2$  it forbids certain pairs of two electric or two magnetic  $p$ -branes, corresponding to the same form ( $F^a$ ,  $a \in \Delta$ ). Restriction 2 is satisfied for  $n_1 = 0$  or when  $d_0 \neq 1, 3$ . For  $n_1 \geq 1$  and  $d_0 = 1, 3$  it forbids certain electro-magnetic pairs, corresponding to the same form.

It was proved in [5] that equations of motion for the model (4.1) and the Bianchi identities:  $d\mathcal{F}^s = 0$ ,  $s \in S_m$ , for fields from (4.3)–(4.13), when Restrictions 1 and 2 are imposed, are equivalent to equations of motion for the  $\sigma$ -model (1.1) with  $(\sigma^A) = (\phi^i, \varphi^\alpha)$ , the index set  $S$  from (4.11), target space metric

$$(\hat{G}_{AB}) = \begin{pmatrix} G_{ij} & 0 \\ 0 & h_{\alpha\beta} \end{pmatrix}, \quad (4.18)$$

with

$$G_{ij} = d_i \delta_{ij} + \frac{d_i d_j}{d_0 - 2}, \quad (4.19)$$

the potential

$$V = -\frac{1}{2} \sum_{i=1}^n \xi_i d_i e^{-2\phi^i + 2\gamma_0(\phi)}, \quad (4.20)$$

vectors

$$(U_A^s) = (d_i \delta_{iI_s}, -\chi_s \lambda_{\alpha a_s}), \quad (4.21)$$

where  $s = (a_s, v_s, I_s)$ ,  $\chi_s = +1, -1$  for  $v_s = e, m$  respectively

$$\delta_{iI} = \sum_{j \in I} \delta_{ij} \quad (4.22)$$

is the indicator of  $i$  belonging to  $I$ :  $\delta_{iI} = 1$  for  $i \in I$  and  $\delta_{iI} = 0$  otherwise; and

$$\varepsilon_s = (-\varepsilon[g])^{(1-\chi_s)/2} \varepsilon(I_s) \theta_{a_s}, \quad (4.23)$$

$s \in S$ ,  $\varepsilon[g] \equiv \text{sign det}(g_{MN})$ . More explicitly (4.23) reads

$$\varepsilon_s = \varepsilon(I_s) \theta_{a_s}, \quad v_s = e \quad (4.24)$$

$$\varepsilon_s = -\varepsilon[g] \varepsilon(I_s) \theta_{a_s} \quad v_s = m. \quad (4.25)$$

The scalar products (1.4) for vectors  $U^s$  were calculated in [5]

$$(U^s, U^{s'}) = d(I_s \cap I_{s'}) + \frac{d(I_s)d(I_{s'})}{2-D} + \chi_s \chi_{s'} \lambda_{\alpha a_s} \lambda_{\beta a_{s'}} h^{\alpha\beta} \equiv B^{ss'}, \quad (4.26)$$

where  $(h^{\alpha\beta}) = (h_{\alpha\beta})^{-1}$ ;  $s = (a_s, v_s, I_s)$  and  $s' = (a_{s'}, v_{s'}, I_{s'})$  belong to  $S$ .

## 4.2 Exact solutions with Ricci-flat spaces

Here we consider the case of Ricci-flat internal spaces, i.e.  $\xi_i = 0$ ,  $i = 1, \dots, n$ , in (4.4). The potential (4.20) is trivial in this case and we may apply the results from Sect.2 to our multidimensional model (4.1), when vectors  $(U^s, s \in S)$  obey the block-orthogonal decomposition (1.6), (1.7) with scalar products defined in (4.26). The solution reads:

$$g = U \left\{ g^0 + \sum_{i=1}^n U_i g^i \right\}, \quad (4.27)$$

$$U = \left( \prod_{s \in S} H_s^{2d(I_s) \varepsilon_s \nu_s^2} \right)^{1/(2-D)}, \quad (4.28)$$

$$U_i = \prod_{s \in S} H_s^{2\varepsilon_s \nu_s^2 \delta_{iI_s}}, \quad (4.29)$$

$$\text{Ric}[g^0] = \text{Ric}[g^1] = \dots = \text{Ric}[g^n] = 0, \quad (4.30)$$

$$\varphi^\alpha = - \sum_{s \in S} \lambda_{a_s}^\alpha \chi_s \varepsilon_s \nu_s^2 \ln H_s, \quad (4.31)$$

$$F^a = \sum_{s \in S} \mathcal{F}^s \delta_{a_s}^a, \quad (4.32)$$

where

$$\mathcal{F}^s = \nu_s dH_s^{-1} \wedge \tau(I_s), \text{ for } v_s = e, \quad (4.33)$$

$$\mathcal{F}^s = \nu_s (*_0 dH_s) \wedge \tau(\bar{I}_s), \text{ for } v_s = m, \quad (4.34)$$

$H_s$  are harmonic functions on  $(M_0, g^0)$  coinciding inside blocks of matrix  $(B^{ss'})$  from (4.26) ( $H_s = H_{s'}$ ,  $s, s' \in S_j$ ,  $j = 1, \dots, k$ , see (1.6), (1.7)) and relations

$$\sum_{s' \in S} B^{ss'} \varepsilon_{s'} \nu_{s'}^2 = -1 \quad (4.35)$$

on the matrix  $(B^{ss'})$ , parameters  $\varepsilon_s$  (4.23) and  $\nu_s$  are imposed,  $s \in S$ ,  $i = 1, \dots, n$ ;  $\alpha = 1, \dots, l$ . Here  $\lambda_a^\alpha = h^{\alpha\beta} \lambda_{\beta a}$ ,  $*_0 = *[g^0]$  is the Hodge operator on  $(M_0, g^0)$  and  $\bar{I}$  is defined in (4.17).

In deriving the solution we used as in [5] the relations for contravariant components of  $U^s$ -vectors:

$$U^{si} = \delta_{iI_s} - \frac{d(I_s)}{D-2}, \quad U^{s\alpha} = -\chi_s \lambda_{a_s}^\alpha, \quad (4.36)$$

$$s = (a_s, v_s, I_s).$$

Thus, we obtained the generalization of the solutions from [5] to the block-orthogonal case (here we eliminate the misprint with sign in eq. (5.19) in [5]).

**Remark 2.** The solution is also valid for  $d_0 = 2$ , if Restriction 2 is replaced by Restriction 2\* . It may be proved using a more general form of the sigma-model representation (see Remark 2 in [5]).

**Restriction 2\*** (for  $d_0 = 2$ ). For any  $a \in \Delta$  there are no  $I \in \Omega_{a,m}$ ,  $J \in \Omega_{a,e}$  such that  $\bar{I} = J$  and for  $n_1 \geq 2$ ,  $i, j \in w_1$ ,  $i \neq j$ , there are no  $I \in \Omega_{a,m}$ ,  $J \in \Omega_{a,e}$  such that  $i \in I$ ,  $j \in \bar{J}$ ,  $I \setminus \{i\} = \bar{J} \setminus \{j\}$ .

### 4.3 Behaviour of the Kretschmann scalar and horizon

Let  $M_0 = \mathbf{R}^{d_0}$ ,  $d_0 > 2$  and  $g^0 = \delta_{\mu\nu} dx^\mu \otimes dx^\nu$ . For

$$H_s = 1 + \sum_{b \in X_s} \frac{q_{sb}}{|x - b|^{d_0-2}}, \quad (4.37)$$

where  $X_s$  is finite non-empty subset  $X_s \subset M_0$ ,  $s \in S$ , all  $q_{sb} > 0$ , and  $X_s = X_{s'}$ ,  $q_{sb} = q_{s'b}$  for  $b \in X_s = X_{s'}$ ,  $s, s' \in S_j$ ,  $j = 1, \dots, k$ . The harmonic functions (4.37) are defined in domain  $M_0 \setminus X$ ,  $X = \bigcup_{s \in S} X_s$ , and generate the solutions (4.27)–(4.35).

Denote  $S(b) \equiv \{s \in S \mid b \in X_s\}$ . We also put  $M_i = \mathbf{R}^{d_i}$ ,  $R[g^i] = \mathcal{K}[g^i] = 0$ ,  $i = 1, \dots, n$ , where

$$\mathcal{K}[g] \equiv R_{MNPQ}[g] R^{MNPQ}[g] \quad (4.38)$$

is the Kretschmann scalar (or Riemann tensor squared). Then for the metric (4.27) we obtain

$$\mathcal{K}[g] = \frac{C' + o(1)}{U^2 |x - b|^4} = [C + o(1)] |x - b|^{4(d_0-2)\eta(b)} \quad (4.39)$$

for  $x \rightarrow b \in X$ , where

$$\eta(b) \equiv \sum_{s \in S(b)} (-\varepsilon_s) \nu_s^2 \frac{d(I_s)}{D-2} - \frac{1}{d_0-2}, \quad (4.40)$$

and  $C = C(b) \geq 0$  ( $C = \text{const}$ ). The relation for  $C = C(b)$  is given in Appendix. Relation (4.39) may be obtained using the relation for  $\mathcal{K}[g]$  in Appendix of Ref. [47]. In what follows we consider non-exceptional  $b \in X$  defined by relations  $C = C(b) > 0$ .

**Remark 3.** It follows from the Appendix that an exceptional point  $b \in X$ , defined by relation  $C = C(b) = 0$ , appears iff (if and only if)

$$U(x) \sim c |x - b|^{-2\alpha}, \quad U(x) U_i(x) \sim c_i, \quad (4.41)$$

for  $x \rightarrow b$ , where  $\alpha = 0, 2$  and  $c, c_i \neq 0$  are constants,  $i = 1, \dots, n$ .

Due to (4.39) the metric (4.27) has no curvature singularity when  $x \rightarrow b \in X$ ,  $C(b) > 0$ , iff

$$\eta(b) \geq 0. \quad (4.42)$$

From (4.40) we see that the metric (4.27) is regular at a “point”  $b \in X$  for  $\varepsilon_s = -1$  and large enough values of  $\nu_s^2$ ,  $s \in S(b)$ . For  $\varepsilon_s = +1$ ,  $s \in S(b)$ , we have a curvature singularity at non-exceptional point  $b \in X$ .

Now we consider a special case:  $d_1 = 1$ ,  $g^1 = -dt \otimes dt$ . In this case we have a horizon, when  $x \rightarrow b \in X$ , iff

$$\xi(b) \equiv \sum_{s \in S(b)} (-\varepsilon_s) \nu_s^2 \delta_{1I_s} - \frac{1}{d_0 - 2} \geq 0. \quad (4.43)$$

This relation follows from the requirement of infinite time propagation of light to  $b \in X$ . If  $\varepsilon_s = -1$ ,  $1 \in I_s$  for all  $s \in S(b)$ , we get

$$\eta(b) < \xi(b), \quad (4.44)$$

$b \in X$ . This follows from the inequalities  $d(I_s) < D - 2$  ( $d_0 > 2$ ).

We note that  $g_{tt} \rightarrow 0$  for  $x \rightarrow b \in X$ , if (4.44) is satisfied. This follows from the relation

$$g_{tt} \sim \text{const} |x - b|^{2(d_0 - 2)(\xi(b) - \eta(b))}, \quad (4.45)$$

$x \rightarrow b$ .

**Remark 4.** Due to relations (4.41) and (4.45) the point  $b \in X$  is non-exceptional if  $g^1 = -dt \otimes dt$  and  $1 \in I_s$ ,  $\varepsilon_s = -1$  for all  $s \in S(b)$ .

Thus, for the metric (4.27) with  $H_s$  from (4.37) there are two dimensionless indicators at the non-exceptional point  $b \in X$ : a) horizon indicator  $\xi(b)$  (corresponding to time  $t$ ) and b) curvature singularity indicator  $\eta(b)$ . These indicators define (for our assumptions) the existence of a horizon and the singularity of the Kretschmann scalar (when  $(M_i, g^i)$  are flat,  $i = 1, \dots, n$ ) at non-exceptional  $b \in X$ .

## 4.4 Generalized MP solutions

Here we consider special black hole solutions for the model (4.1) with all  $\theta_a = 1$ ,  $a \in \Delta$ , when the signature of the metric  $g$  is  $(-1, +1, \dots, +1)$ . From the results of the previous subsections we get the following solution (with all  $\varepsilon_s = \varepsilon(I_s) = -1$ )

$$g = \left( \prod_{s \in S} H_s^{2d(I_s)\nu_s^2} \right)^{1/(D-2)} \left\{ \sum_{\mu=1}^{d_0} dx^\mu \otimes dx^\mu - \left( \prod_{s \in S} H_s^{-2\nu_s^2} \right) dt \otimes dt + \sum_{i=2}^n \left( \prod_{s \in S} H_s^{-2\nu_s^2 \delta_{iI_s}} \right) g^i \right\}, \quad (4.46)$$

$$\varphi^\alpha = \sum_{s \in S} \lambda_{a_s}^\alpha \chi_s \nu_s^2 \ln H_s, \quad (4.47)$$

$$F^a = \sum_{s \in S_e} \delta_{a_s}^a \nu_s dH_s^{-1} \wedge \tau(I_s) + \sum_{s \in S_m} \delta_{a_s}^a \nu_s (*_0 dH_s) \wedge \tau(\bar{I}_s), \quad (4.48)$$

where  $(M_i, g^i)$  are flat Euclidean spaces,  $i = 2, \dots, n$ ,

$$1 \in I_s, \quad (4.49)$$

$s \in S$ , i.e. all branes have a common time submanifold  $M_1 = \mathbf{R}$ , the harmonic functions  $H_s$  (4.37) are coinciding inside blocks, i.e.  $H_s = H_{s'}$ ,  $s, s' \in S_j$ ,  $j = 1, \dots, k$ , parameters  $\nu_s$  satisfy the relations

$$\sum_{s' \in S} B^{ss'} \nu_{s'}^2 = 1 \quad (4.50)$$

with the matrix  $(B^{ss'})$  from (4.26),  $*_0 = *[g^0]$  is the Hodge operator on  $(M_0 = \mathbf{R}^{d_0}, g^0 = \sum_{\mu=1}^{d_0} dx^\mu \otimes dx^\mu)$ ,  $\bar{I}$  is defined in (4.17),  $\theta_a = 1$  for all  $a \in \Delta$ , and

$$\eta(b) = \sum_{s \in S(b)} \nu_s^2 \frac{d(I_s)}{D-2} - \frac{1}{d_0-2} \geq 0, \quad (4.51)$$

$b \in X$ . This solution describes a set of extreme  $p$ -brane black holes with horizons at  $b \in X$ . The Riemann tensor squared has a finite limit at any  $b \in X$ .

Calculation of the Hawking temperature corresponding to  $b \in X$  using standard formula (see, for example, [48, 32]) gives us

$$T_H(b) = 0, \quad (4.52)$$

for any  $b \in X$  satisfying  $\xi(b) > 0$ .

**MP solution.** The standard 4-dimensional Majumdar-Papapetrou solution [1] in our notations reads

$$g = H^2 g^0 - H^{-2} dt \otimes dt, \quad (4.53)$$

$$F = \nu dH^{-1} \wedge dt, \quad (4.54)$$

where  $\nu^2 = 2$ ,  $g^0 = \sum_{i=1}^3 dx^i \otimes dx^i$  and  $H$  is a harmonic function. We have one electric 0-brane (point) “attached” to the time manifold;  $d(I_s) = 1$ ,  $\varepsilon_s = -1$  and  $(U^s, U^s) = 1/2$ . In this case (e.g. for extremal Reissner-Nordström black hole) we get

$$\eta(b) = 0, \quad \xi(b) = 1, \quad (4.55)$$

and  $T_H(b) = 0$ ,  $b \in X$ .

**D = 11 supergravity.** Let us consider the action (4.1) without scalar fields and with the only one form  $F^4$  of rank 4. This action coincides with the truncated (i.e. without Chern-Simons term) bosonic part of  $D = 11$  supergravity action. In this case there are a lot of  $p$ -brane MP-type (multicenter extremal black hole) solutions with orthogonal intersection rules [11]-[16], e.g. (i) solution with one electric 2-brane ( $d(I_s) = 3$ ) and  $d_0 = 8$ ; (ii) solution with one magnetic 5-brane ( $d(I_s) = 6$ ) and  $d_0 = 5$ ; (iii) solution with one electric 2-brane and one magnetic 5-brane ( $d(I_{s_1} \cap I_{s_2}) = 2$ ) and  $d_0 = 4$ ; (iv) solution with two electric 2-branes ( $d(I_{s_1} \cap I_{s_2}) = 1$ ) and  $d_0 = 5$ ; (v) solution with two magnetic 5-branes ( $d(I_{s_1} \cap I_{s_2}) = 4$ ) and  $d_0 = 3$ . In the examples (iii)-(v) the harmonic functions  $H_{s_1}$  and  $H_{s_2}$  from (4.37) should have the coinciding sets of poles, i.e.  $X_{s_1} = X_{s_2}$ , to maintain the relation (4.51). The addition of the Chern-Simons term does not destroy these solutions. In all these examples  $\eta(b) = 0$ ,  $b \in X_s$  and  $\nu_s^2 = 1/2$ ,  $s \in S$ .

## 4.5 Non-extremal $p$ -brane black hole solutions

There exists a non-extremal generalization of the solution from the previous subsection, when all the poles in all harmonic functions are coinciding, say  $b = 0$ ,  $b \in X$ .

The metric and scalar fields for this solution read

$$g = \left( \prod_{s \in S} H_s^{2d(I_s)\nu_s^2/(D-2)} \right) \left\{ \frac{dr \otimes dr}{1 - 2\mu/r^{\bar{d}}} + r^2 d\Omega_{d_0-1}^2 \right. \\ \left. - \left( \prod_{s \in S} H_s^{-2\nu_s^2} \right) \left( 1 - \frac{2\mu}{r^{\bar{d}}} \right) dt \otimes dt + \sum_{i=2}^n \left( \prod_{s \in S} H_s^{-2\delta_{iI_s}\nu_s^2} \right) g^i \right\}, \quad (4.56)$$

$$\varphi^\alpha = \sum_{s \in S} \nu_s^2 \chi_s \lambda_{a_s}^\alpha \ln H_s, \quad (4.57)$$

where

$$H_s = 1 + \frac{q_s}{r^{\bar{d}}}, \quad (4.58)$$

$s \in S$ ,  $\bar{d} = d_0 - 2$ ,  $\mu > 0$ , parameters  $\nu_s$  satisfy the relations (4.50) with the matrix  $(B^{ss'})$  from (4.26), and parameters  $q_s > 0$  coincide inside blocks i.e.  $q_s = q_{s'}$ ,  $s, s' \in S_j$ ,  $j = 1, \dots, k$ . Here  $(M_i, g^i)$  are Ricci-flat Euclidean spaces,  $i = 2, \dots, n$ , and  $d\Omega_{d_0-1}^2$  is canonical metric on the  $(d_0 - 1)$ -dimensional sphere  $S^{d_0-1}$ . The fields of forms are given by (4.32)-(4.34) with

$$\Phi^s = \frac{\nu_s}{H'_s}, \quad (4.59)$$

$$H'_s = \left( 1 - \frac{p_s}{H_s r^{\bar{d}}} \right)^{-1} = 1 + \frac{p_s}{r^{\bar{d}} + q_s - p_s}, \quad (4.60)$$

$$p_s = \sqrt{q_s(q_s + 2\mu)}, \quad (4.61)$$

$s \in S$ .

The solution describes non-extremal charged intersecting  $p$ -brane black hole with the horizon at  $r^{\bar{d}} = 2\mu$ . In the limit  $\mu \rightarrow +0$  this solution coincides with the 1-center extremal black hole solution from the previous subsection. Here we generalize the solution from [33], where all  $\nu_s^2$  are coinciding inside blocks. In our case the only restriction on  $\nu_s$  is the relation (4.50). For orthogonal intersections it agrees with the solutions from [24]-[27] ( $d_1 = \dots = d_n = 1$ ) and [32, 35].

The Hawking temperature corresponding to this non-extremal solution is

$$T_H(0, \mu) = \frac{\bar{d}}{4\pi(2\mu)^{1/\bar{d}}} \prod_{s \in S} \left( \frac{2\mu}{2\mu + q_s} \right)^{\nu_s^2}. \quad (4.62)$$

For  $\mu \rightarrow +0$  we get (in agreement with (4.52))  $T_H(0, \mu) \rightarrow 0$  for the extremal black hole configurations satisfying

$$\xi(0) = \sum_{s \in S} \nu_s^2 - \bar{d}^{-1} > 0. \quad (4.63)$$

## 5 Intersection rules and some examples

### 5.1 Intersection rules

From orthogonality relation (1.7) and (4.26) we get

$$d(I_s \cap I_{s'}) = \Delta(s, s') \quad (5.1)$$

where  $s \in S_i$ ,  $s' \in S_j$ ,  $i \neq j$  and

$$\Delta(s, s') \equiv \frac{d(I_s)d(I_{s'})}{D-2} - \chi_s \chi_{s'} \lambda_{a_s} \cdot \lambda_{a_{s'}}. \quad (5.2)$$

Here  $\lambda \cdot \lambda' \equiv h^{\alpha\beta} \lambda_\alpha \lambda'_\beta$ . Let

$$N(a, b) \equiv \frac{(n_a - 1)(n_b - 1)}{D - 2} - \lambda_a \cdot \lambda_b, \quad (5.3)$$

$a, b \in \Delta$ . The matrix (5.3) is called the fundamental matrix of the model (4.1) [35]. For  $s_1, s_2 \in S$ ,  $s_1 \neq s_2$ , the  $\Delta$ -symbol (5.2) may be expressed by means of the fundamental matrix [35]

$$\Delta(s_1, s_2) = \bar{D} \bar{\chi}_{s_1} \bar{\chi}_{s_2} + \bar{n}_{a_{s_1}} \chi_{s_1} \bar{\chi}_{s_2} + \bar{n}_{a_{s_2}} \chi_{s_2} \bar{\chi}_{s_1} + N(a_{s_1}, a_{s_2}) \chi_{s_1} \chi_{s_2}, \quad (5.4)$$

where  $\bar{D} = D - 2$ ,  $\bar{n}_a = n_a - 1$ ,  $\bar{\chi}_s = \frac{1}{2}(1 - \chi_s)$ . More explicitly (5.4) reads

$$\Delta(s_1, s_2) = N(a_{s_1}, a_{s_2}), \quad v_{s_1} = v_{s_2} = e; \quad (5.5)$$

$$\Delta(s_1, s_2) = \bar{n}_{a_{s_1}} - N(a_{s_1}, a_{s_2}), \quad v_{s_1} = e, \quad v_{s_2} = m; \quad (5.6)$$

$$\Delta(s_1, s_2) = \bar{D} - \bar{n}_{a_{s_1}} - \bar{n}_{a_{s_2}} + N(a_{s_1}, a_{s_2}), \quad v_{s_1} = v_{s_2} = m. \quad (5.7)$$

This follows from the relations

$$d(I_s) = \bar{D} \bar{\chi}_s + \bar{n}_{a_s} \chi_s, \quad (5.8)$$

equivalent to (4.12). Let

$$K(a) \equiv n_a - 1 - N(a, a) = \frac{(n_a - 1)(D - n_a - 1)}{D - 2} + \lambda_a \cdot \lambda_a, \quad (5.9)$$

$a \in \Delta$ .

The parameters (5.9) play a rather important role in supergravity theories, since they are preserved under Kaluza-Klein reduction [21] and define the norms of  $U^s$  vectors:

$$(U^s, U^s) = K(a_s), \quad (5.10)$$

$s \in S$ .

Here we put  $K(a) \neq 0$ ,  $a \in \Delta$ . Then, we obtain the general intersection rule formulas

$$d(I_{s_1} \cap I_{s_2}) = \Delta(s_1, s_2) + \frac{1}{2} K(a_{s_2}) A^{s_1 s_2} \quad (5.11)$$

$s_1 \neq s_2$ , where  $(A^{s_1 s_2})$  is the quasi-Cartan matrix (3.2) (see also (6.32) from [35]).

## 5.2 $B_D$ -models and examples of solutions

Now we consider some examples of solutions from Sect. 4. These examples will be demonstrated for the so-called  $B_D$ -models. Action of the  $B_D$ -model reads [35]

$$S_D = \int d^D z \sqrt{|g|} \left\{ R[g] + g^{MN} \partial_M \vec{\varphi} \partial_N \vec{\varphi} - \sum_{a=4}^{D-7} \frac{1}{a!} \exp[2\vec{\lambda}_a \vec{\varphi}] (F^a)^2 \right\}, \quad (5.12)$$

where  $\vec{\varphi} = (\varphi^1, \dots, \varphi^l) \in \mathbf{R}^l$ ,  $\vec{\lambda}_a = (\lambda_{a1}, \dots, \lambda_{al}) \in \mathbf{R}^l$ ,  $l = D - 11$ ,  $\text{rank } F^a = a$ ,  $a = 4, \dots, D - 7$ . Here vectors  $\vec{\lambda}_a$  satisfy the relations

$$\vec{\lambda}_a \vec{\lambda}_b = N(a, b) - \frac{(a - 1)(b - 1)}{D - 2}, \quad (5.13)$$

$$N(a, b) = \min(a, b) - 3, \quad (5.14)$$

$a, b = 4, \dots, D-7$ .

The vectors  $\vec{\lambda}_a$  are linearly dependent

$$\vec{\lambda}_{D-7} = -2\vec{\lambda}_4. \quad (5.15)$$

For  $D > 11$  vectors  $\vec{\lambda}_4, \dots, \vec{\lambda}_{D-8}$  are linearly independent.

The model (5.12) contains  $l$  scalar fields with a negative kinetic term (i.e.  $h_{\alpha\beta} = -\delta_{\alpha\beta}$  in (4.1)) coupled to  $(l+1)$  forms. For  $D = 11$  ( $l = 0$ ) the model (5.12) coincides with a truncated (without Chern-Simons term) bosonic sector of  $D = 11$  supergravity. For  $D = 12$  ( $l = 1$ ) (5.12) coincides with truncated  $D = 12$  model from [30] (see also [5]).

The matrix (5.14) is the fundamental matrix of the  $B_D$ -model.

For  $p$ -brane worldsheets we have the following dimensions (see [35])

$$d(I) = 3, \dots, D-8, \quad I \in \Omega_{a,e}, \quad (5.16)$$

$$d(I) = D-5, \dots, 6, \quad I \in \Omega_{a,m}. \quad (5.17)$$

Thus, there are  $(l+1)$  electric and  $(l+1)$  magnetic  $p$ -branes,  $p = d(I) - 1$ . For  $B_D$ -model all  $K(a) = 2$ .

Now we are interested in the one-block solutions, where  $A = (A^{s_1 s_2})$  in (3.2) is an irreducible Cartan matrix either of a simple finite dimensional Lie algebra or of a simple hyperbolic KM algebra.

Since  $K(a) = 2$ , we rewrite (5.11) as the following:

$$d(I_{s_1} \cap I_{s_2}) = \Delta(s_1, s_2) + A^{s_1 s_2}, \quad (5.18)$$

$s_1 \neq s_2$ , and get  $A^{s_1 s_2} = A^{s_2 s_1}$ , i.e. the Cartan matrix is symmetric.

### 5.2.1 Finite dimensional Lie algebras

Here we put all  $\theta_a = 1$ ,  $a \in \Delta$ . From  $\varepsilon_s = -1$  we get  $\varepsilon(I_s) = -1$  for the electric "brane" and  $\varepsilon(I_s) = \varepsilon[g]$  for the magnetic one. In a finite dimensional case we are led to the so-called simply laced or  $A - D - E$  Lie algebras. The intersection rules are totally defined by the corresponding Dynkin diagram:  $d(I_{s_1} \cap I_{s_2}) = \Delta(s_1, s_2) - 1$ , when the vertices corresponding to  $s_1$  and  $s_2$  are connected by a line and  $d(I_{s_1} \cap I_{s_2}) = \Delta(s_1, s_2)$  otherwise (since in  $A - D - E$  case  $A^{s_1 s_2} = 0, -1$ ,  $s_1 \neq s_2$ ).

**Example for  $A_2$ .** Let us consider  $B_D$ -model,  $D \geq 11$ ,  $a \in \{4, \dots, D-7\}$ ,  $g^3 = -dt \otimes dt$ ,  $d_1 = a - 2$ ,  $d_2 = D - 2 - a$ ,  $d_0 = 3$  and  $g^0, g^1, g^2$  are Ricci-flat. The  $A_2$ -solution describing a dyon configuration with electric  $d_1$ -brane and magnetic  $d_2$ -brane, corresponding to  $F^a$ -form and intersecting in 1-dimensional time manifold reads:

$$g = H^2 g^0 - H^{-2} dt \otimes dt + g^1 + g^2, \quad (5.19)$$

$$F^a = \nu_1 dH^{-1} \wedge dt \wedge \tau_1 + \nu_2 (*_0 dH) \wedge \tau_1, \quad (5.20)$$

$\vec{\varphi} = 0$ ,  $H$  is the harmonic function on  $(M_0, g^0)$  and  $\nu_1^2 = \nu_2^2 = 1$ . For  $D = 11$  we have  $a = 4$  and  $d_1 = 2$ ,  $d_2 = 5$ . For  $D = 12$  we have two possibilities: a)  $a = 4$ ,  $d_1 = 2$ ,  $d_2 = 6$ ; b)  $a = 5$ ,  $d_1 = 3$ ,  $d_2 = 5$ . The signature restrictions on  $g^1$  and  $g^2$  are the following:  $\varepsilon(1) = +1$ ,  $\varepsilon(2) = -\varepsilon[g]$ . They are satisfied when  $g^0$  and  $g^1$  are Euclidean metrics. The 4-dimensional section of (5.19) coincides for the flat Euclidean  $g^0$  with the MP solution (4.53), (4.54). Here the "indicators" of the solution coincide with MP ones (4.55).

Now, we list  $A_2$  intersection rules, for  $D = 11, 12$ . Here

$$A = B = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}. \quad (5.21)$$

**D = 11 :**

$$3 \cap 3 = 0, \quad 3 \cap 6 = 1, \quad 6 \cap 6 = 3. \quad (5.22)$$

**D = 12 :**

$$\begin{aligned} 3 \cap 3 = 3 \cap 4 = 0, \quad 3 \cap 6 = 3 \cap 7 = 4 \cap 4 = 4 \cap 6 = 1, \\ 4 \cap 7 = 2, \quad 6 \cap 6 = 6 \cap 7 = 3, \quad 7 \cap 7 = 4. \end{aligned} \quad (5.23)$$

Here and in what follows we denote  $n_1 \cap n_2 = n \Leftrightarrow (d(I_1) = n_1, d(I_2) = n_2, d(I_1 \cap I_2) = n)$ .

**Remark 5.** The appearance of Lie algebras it is not accident here. It was shown in [35] that if  $p$ -branes have intersection rules (5.11) governed by a Cartan matrix of some (semisimple) Lie algebra, then equations of motion for the problems of cosmology and spherical symmetry may be reduced to integrable Euclidean Toda lattice equations corresponding to this Lie algebra. (For certain examples of solutions see, for example, [53, 54, 55, 56].) Thus, for intersections related to Lie algebras we may find general spherically symmetric solutions which contain one-center MP-type solutions (e.g. extremal black hole configurations) as special case.

### 5.2.2 Hyperbolic algebras

In hyperbolic case all  $\varepsilon_s = +1$ ,  $s \in S_1 = S$  and, hence, corresponding solutions with  $H_s$  from (4.37) are singular at  $b \in X$  (in this case  $b$  is non-exceptional).

**Example with  $H_2(q, q)$ .** For the Cartan matrix (3.13) with  $q_1 = q_2 = q \geq 3$ , we obtain

$$\nu_s^2 = (q - 2)^{-1} = 1, \frac{1}{2}, \frac{1}{3}, \dots \quad (5.24)$$

for  $q = 3, 4, 5, \dots$ ;  $s = 1, 2$ . An example of the solution for  $d_0 = 3$  with two electric  $p$ -branes,  $p = d_1, d_2$ , corresponding to  $F^a$  and  $F^b$  fields and intersecting in time manifold, is the following:

$$g = H^{-2/(q-2)} g^0 - H^{2/(q-2)} dt \otimes dt + g^1 + g^2, \quad (5.25)$$

$$F = \nu_1 dH^{-1} \wedge dt \wedge \tau_1 + \nu_2 dH^{-1} \wedge dt \wedge \tau_2, \quad (5.26)$$

$$\vec{\varphi} = -(\vec{\lambda}_a + \vec{\lambda}_b)(q - 2)^{-1} \ln H \quad (5.27)$$

where  $d_1 = a - 2$ ,  $a = q + 4$ ,  $b \geq a$ ,  $d_2 = b - 2$ ,  $d_0 = 3$ ,  $D = a + b$ . Here  $F = F^a + F^b$  for  $a < b$  and  $F = F^a$  for  $a = b$ . The signature restrictions are :  $\varepsilon(1) = \varepsilon(2) = -1$ . Thus, the space-time  $(M, g)$  should contain at least three time directions. The minimal  $D$  is 14. For  $D = 14$  we get  $a = b = 7$ ,  $d_1 = d_2 = 6$ ,  $q = 3$ . In this case  $6 \cap 6 = 1$ .

### 5.2.3 Affine (forbidden) case

Affine Cartan matrices do not arise in our solutions. This means that some configurations are forbidden. Let us consider  $A_1^{(1)}$  affine KM algebra with the Cartan matrix

$$A = \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}. \quad (5.28)$$

For  $D = 11$  the intersections:  $3 \cap 6 = 0$ ,  $6 \cap 6 = 2$ , corresponding to the  $A$ -matrix (5.28), are forbidden. For  $D = 12$  we get forbidden intersections:  $3 \cap 6 = 3 \cap 7 = 4 \cap 4 = 4 \cap 6 = 0$ ,  $4 \cap 7 = 1$ ,  $6 \cap 6 = 6 \cap 7 = 2$ ,  $7 \cap 7 = 3$ .

**Remark 6.** Recently some new solutions in the affine case were obtained [49]. (These solutions contain as a special case a solution in  $D = 11$  supergravity from [16] with  $6 \cap 6 = 2$ .) The solutions from [49] use some modified ansatz for fields of forms and do not belong to our scheme. This indicates that the sigma-model considered in this paper is not the most general one describing possible  $p$ -brane configurations. There are at least three possible ways of generalization: (i) “non-block-diagonal” metric instead of the “block-diagonal” metric (4.3) may be considered (the first step in this direction was done in [34]); (ii) a bimetric sigma-model defined on the product of two base spaces  $(M_{01}, g^{01})$  and  $(M_{02}, g^{02})$  instead of  $(M_0, g^0)$  with two sorts of  $p$ -branes governed by functions on  $M_{01}$  and  $M_{02}$  respectively may be constructed; (iii) a sigma-model describing the action (4.1) with Chern-Simons terms added may be also considered. Solution from [49] seems to belong to the case (ii). We note also that our solutions in the special case of  $D = 10, 11$  supergravities are different from non-marginal bound state solutions (see, for example, [27, 50, 51] and references therein) although the rules for binary intersections may look similar. These solutions probably need some extensions of the sigma-model along the lines (i) and (iii).

## 5.2.4 Other possibilities

We note that it is not obvious for quasi-Cartan matrix  $A$  to be a Cartan one. Let us consider the example with

$$A = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}. \quad (5.29)$$

For  $D = 11$  we obtain (non-Lie) dyon with

$$3 \cap 6 = 3 \quad (5.30)$$

and  $\varepsilon_s = -1$ ,  $\nu_s^2 = 1/3$ ,  $s = 1, 2$  (see also [33]). Two other intersections:  $3 \cap 3 = 2$ ,  $6 \cap 6 = 5$ , corresponding to  $A$  are forbidden by Restriction 1 (this restriction follows from the block-diagonal form of metric  $g$  and may be weakened for non-block-diagonal ansatz from [34]). For  $d_0 = 5$  the solution with  $d_1 = d_2 = 3$  reads

$$\begin{aligned} g &= H^{2/3} g^0 + H^{-2/3} g^1 + g^2, \\ F^4 &= \nu_1 dH^{-1} \wedge \tau_1 + \nu_2 *_0 dH \end{aligned} \quad (5.31)$$

with  $H$  from (4.37) and indicators  $\xi(b) = 1/3$ ,  $\eta(b) = 0$ ,  $b \in X$ , i.e. any point  $a \in X$  corresponds to the horizon without a curvature singularity. Here we assume that  $M_1 = \mathbf{R} \times M'_1$ , where  $\mathbf{R}$  is the time manifold.

## 6 Conclusions

Thus, here, like in [3]–[6], we considered the  $\sigma$ -model of a  $p$ -brane origin. We obtained new block-orthogonal exact solutions (2.4)–(2.7) governed by a set of harmonic functions. These solutions crucially depend on the matrix of scalar products  $(U^s, U^{s'})$ ,  $s, s' \in S$ , and parameters  $\varepsilon_s = \pm 1$ ,  $s \in S$  (see (1.8)). In Sect. 3 we analyzed three possibilities, when quasi-Cartan matrix (3.2) coincides

with the Cartan matrix of a (simple): a) finite-dimensional Lie algebra; b) hyperbolic KM algebra; c) affine KM algebra. It was shown that the last possibility does not appear in our solutions, and all  $\varepsilon_s = -1$  in the case a) and all  $\varepsilon_s = +1$  in the case b). In Sect. 4 we applied this method for obtaining new MP type solutions in multidimensional gravity with fields of forms and scalar fields (see subsect. 4.2 and 4.4). In subsect. 4.3 for  $d_0 > 2$  and harmonic functions from (4.37) the indicators  $\eta(b)$  and  $\xi(b)$  were introduced. These indicators describe under certain assumptions the existence of a curvature singularity and a horizon for  $x \rightarrow b$ . In Sect. 5 intersection rules for fixed  $A$ -matrix are written (see (5.11)) and some examples of solutions and intersection rules for a chain of  $B_D$ -models,  $D = 11, 12, \dots$ , were suggested. Among the examples there are  $A_2$ , hyperbolic, affine (forbidden) intersections and “non-Cartan” ones. We note that the solutions obtained here may be also generalized to the case when some non-Ricci-flat internal spaces are added to  $M$  as it was done in [5].

## 7 Appendix

Here we present the relation for the parameter  $C = C(b)$ ,  $b \in X$ , from (4.39)

$$C = C_0 + C_1 + C_2, \quad (7.1)$$

$$C_0 = 2(d_0 - 1)(d_0 - 2)\alpha^2(\alpha - 2)^2, \quad (7.2)$$

$$C_1 = 4[(d_0 - 1)\alpha^2 + (\alpha - 1)^2] \sum_{i=1}^n d_i \alpha_i^2, \quad (7.3)$$

$$C_2 = 2\left(\sum_{i=1}^n d_i \alpha_i^2\right)^2 - 2 \sum_{i=1}^n d_i \alpha_i^4, \quad (7.4)$$

where

$$\alpha = \alpha(b) \equiv (d_0 - 2) \sum_{s \in S(b)} (-\varepsilon_s) \nu_s^2 \frac{d(I_s)}{D - 2} = (d_0 - 2)\eta(b) + 1, \quad (7.5)$$

$$\alpha_i = \alpha_i(b) \equiv (d_0 - 2) \sum_{s \in S(b)} (-\varepsilon_s) \nu_s^2 \left[ \delta_{iI_s} - \frac{d(I_s)}{D - 2} \right], \quad (7.6)$$

$i = 1, \dots, n$ .

It follows from definitions (7.1)-(7.4) that  $C \geq 0$  and

$$C = 0 \Leftrightarrow (\alpha = 0, 2, \quad \alpha_i = 0, \quad i = 1, \dots, n). \quad (7.7)$$

Parameter  $C$  appears in the Kretschmann scalar (4.39) for the “1-pole” metric

$$g_* = r^{-2\alpha} [dr \otimes dr + r^2 d\Omega_{d_0-1}^2] + \sum_{i=1}^n r^{2\alpha_i} g^i, \quad (7.8)$$

with  $R[g^i] = \mathcal{K}[g^i] = 0$ ,  $i = 1, \dots, n$ . Using formulas from Appendix of [47], we obtain

$$\mathcal{K}[g_*] = C r^{-4+4\alpha}. \quad (7.9)$$

### Acknowledgments

This work was supported in part by the DFG grant 436 RUS 113/236/O(R), by the Russian Ministry of Science and Technology and Russian Foundation for Basic Research. The authors are grateful to M.A. Grebeniuk, K.A. Bronnikov, S.-W. Kim for useful discussions and V.V. Nikulin for valuable information. One of us (V.D.I) also thanks S.-W. Kim for a kind hospitality during his stay in Ewha Womans University (Seoul, S. Korea).

# References

- [1] S.D. Majumdar, *Phys. Rev.* **72**, 930 (1947);  
A. Papapetrou, *Proc. R. Irish Acad.* **A51**, 191 (1947).
- [2] S.V. Chervon, Non-Linear Fields in General Relativity and Cosmology, Ulyanovsk, 1997 (in Russian).
- [3] V.D. Ivashchuk and V.N. Melnikov, Intersecting p-brane Solutions in Multidimensional Gravity and M-theory, hep-th/9612089; *Gravitation and Cosmology* **2**, No 4, 204 (1996).
- [4] V.D. Ivashchuk and V.N. Melnikov, *Phys. Lett.* **B 403**, 23 (1997).
- [5] V.D. Ivashchuk and V.N. Melnikov, Sigma-model for the Generalized Composite p-branes, hep-th/9705036; *Class. Quantum Grav.* **14** (11), 3001 (1997).
- [6] V.D. Ivashchuk, V.N. Melnikov and M. Rainer, Multidimensional  $\sigma$ -models with Composite Electric p-branes, gr-qc/9705005; *Gravitation and Cosmology* **4**, No. 1 (13), 73 (1998).
- [7] V.A. Berezin, G. Domenech, M.L. Levinas, C.O. Lousto and N.D. Umerz, *Gen. Relativ. Grav.* **21**, 1177 (1989).
- [8] M. Rainer and A. Zhuk, *Phys. Rev.*, **D 54** 6186 (1996).
- [9] V.D. Ivashchuk and V.N. Melnikov, Multidimensional Gravity with Einstein Internal spaces, hep-th/9612054; *Gravitation and Cosmology* **2**, No 3 (7), 177 (1996).
- [10] A. Dabholkar, G. Gibbons, J.A. Harvey, and F. Ruiz Ruiz, *Nucl. Phys.* **B 340**, 33 (1990).
- [11] M.J. Duff and K.S. Stelle, *Phys. Lett.* **B 253**, 113 (1991).
- [12] R. Güven, *Phys. Lett.* **B 276**, 49 (1992); *Phys. Lett.* **B 212**, 277 (1988).
- [13] M.J. Duff, R.R. Khuri and J.X. Lu, *Phys. Rep.* **259**, 213 (1995).
- [14] G. Papadopoulos and P.K. Townsend, *Phys. Lett.* **B 380**, 273 (1996); hep-th/9603087.
- [15] A.A. Tseytlin, *Nucl. Phys.* **B 475**, 149 (1996); hep-th/9604035.
- [16] J.P. Gauntlett, D.A. Kastor, and J. Traschen, *Nucl. Phys.* **B 478**, 544 (1996); hep-th/9604179.
- [17] I.Ya. Aref'eva and A.I. Volovich, *Class. Quantum Grav.* **14** (11), 2991 (1997); hep-th/9611026.
- [18] E. Bergshoeff, M. de Roo, E. Eyras, B. Janssen and J.P. van der Schaar, *Class. Quantum Grav.* **14**, 2757 (1997); hep-th/9612095.
- [19] I.Ya. Aref'eva and O.A. Rytchkov, Incidence Matrix Description of Intersecting p-brane Solutions, *Preprint SMI-25-96*, hep-th/9612236.
- [20] R. Argurio, F. Englert and L. Hourant, *Phys. Lett.* **B 398**, 2991 (1997); hep-th/9701042.
- [21] K.S. Stelle, Lectures on Supergravity p-branes, hep-th/9701088.
- [22] J. P. Gauntlett, Intersecting Branes, hep-th/9705011.
- [23] I.Ya. Aref'eva, M.G. Ivanov and O.A. Rytchkov, Properties of Intersecting p-branes in Various Dimensions, *Preprint SMI-05-97*, hep-th/9702077.
- [24] M. Cvetič and A.A. Tseytlin, *Nucl. Phys.* **B 478**, 181 (1996).

- [25] N. Ohta and T. Shimizu, Non-extreme Black Holes from Intersecting M-branes, hep-th/9701095.
- [26] I.Ya. Aref'eva, M.G. Ivanov and I.V. Volovich, Non-extremal Intersecting p-branes in Various Dimensions, hep-th/9702079; *Phys. Lett. B* **406**, 44 (1997).
- [27] N. Ohta, Intersection Rules for Non-extreme p-branes, hep-th/9702164.
- [28] C. Hull and P. Townsend, *Nucl. Phys. B* **438**, 109 (1995); hep-th/9610167;  
P. Horava and E. Witten, *Nucl. Phys. B* **460**, 506 (1996); hep-th/9510209;  
J.M. Schwarz, Lectures on Superstring and M-theory Dualities, hep-th/9607201;  
M.J. Duff, M-theory (the Theory Formerly Known as Strings), hep-th/9608117.
- [29] C.M. Hull, *Nucl. Phys. B* **468**, 113 (1996);  
C. Vafa, *Nucl. Phys. B* **469**, 403 (1996); hep-th/9602022.
- [30] N. Khvengia, Z. Khvengia, H. Lü and C.N. Pope, Toward Field Theory of F-Theory, hep-th/9703012.
- [31] K.A. Bronnikov, M.A. Grebeniuk, V.D. Ivashchuk and V.N. Melnikov, Integrable Multidimensional Cosmology for Intersecting  $p$ -branes, gr-qc/9709006; *Gravitation and Cosmology* **3**, No 2(10), 105 (1997).
- [32] K.A. Bronnikov, V.D. Ivashchuk and V.N. Melnikov, The Reissner-Nordström Problem for Intersecting Electric and Magnetic  $p$ -branes, gr-qc/9710054; *Gravitation and Cosmology*, **3**, No 3(11), 203 (1997).
- [33] K.A. Bronnikov, Block-orthogonal Brane systems, Black Holes and Wormholes, hep-th/9710207; *Gravitation and Cosmology* **4**, No 1(13), 49 (1998).
- [34] D.V. Gal'tsov and O.A. Rytchkov, Generating Branes via Sigma models, hep-th/9801180.
- [35] V.D. Ivashchuk and V.N. Melnikov, Multidimensional Classical and Quantum Cosmology with Intersecting  $p$ -branes, hep-th/9708157; *J. Math. Phys.*, **39**, 2866 (1998).
- [36] V.D. Ivashchuk, S.-W. Kim and V.N. Melnikov, Hyperbolic Lie Algebra from Intersecting p-branes, hep-th/9802054, submitted to J. Math. Phys.
- [37] M.A. Grebeniuk and V.D. Ivashchuk, Solutions in Sigma Model and Intersecting p-branes Related to the Lie Algebras, hep-th/9805113; submit. to Phys. Lett. B.
- [38] V.D. Ivashchuk and V.N. Melnikov, On Symmetries of Target Space for Sigma-model of  $p$ -brane Origin, Proc. of Cosmion meeting, Moscow, December, 7–12, 1997, World Scientific, Singapore (to be published); hep-th/9804102; *Gravitation and Cosmology* **4**, No 3(15) (1998).
- [39] V.G. Kac, Infinite-dimensional Lie Algebras (Cambridge University Press, Cambridge, 1990).
- [40] J. Fuchs and C. Schweigert, Symmetries, Lie algebras and Representations (Cambridge University Press, Cambridge, 1997).
- [41] V.V. Nikulin, private communication.
- [42] A. Feingold, I.B. Frenkel, *Math. Ann.* **263**, 87 (1983).
- [43] B. Julia, in *Lectures in Applied Mathematics*, **21**, 355 (1985).
- [44] S. Mizoguchi,  $E_{10}$  Symmetry in One Dimensional Supergravity, hep-th/9703160.
- [45] H. Nicolai, *Phys. Lett. B* **276**, 333 (1992).

- [46] V.V. Nikulin, On the Classification of Hyperbolic Root Systems of the Rank Three, Part I, [alg-geom/9711032](#).
- [47] V.D. Ivashchuk and V.N. Melnikov, *Class. and Quant. Grav.* **11**, 1793 (1994).
- [48] R. Wald, General Relativity, (University of Chicago, Chicago, 1984).
- [49] J.D. Edelstein, L. Tataru and R. Tatar, Rules for Localized Overlappings and Intersections of p-branes, [hep-th/9801049](#).
- [50] J.M. Izquiero, N.D. Lambert, G. Papadopoulos and P.K. Townsend, *Nucl. Phys.* **460**, 560 (1996); [hep-th/9508177](#).
- [51] J.G. Russo and A.A. Tseytlin, *Nucl. Phys.* **B 490**, 121 (1997); [hep-th/9611047](#).
- [52] N. Ohta and J.-G. Zhou, Towards the Classification of Non-Marginal Bound States of M-branes and Their Construction Rules, [hep-th/9706153](#).
- [53] H. Lü, C.N. Pope, and K.W. Xu, "Liouville and Toda Solitons in M-theory", [hep-th/9604058](#).
- [54] H. Lü, S. Mukherji, C.N. Pope and K.-W. Xu, "Cosmological Solutions in String Theories", [hep-th/9610107](#).
- [55] H. Lü, J. Maharana, S. Mukherji and C.N. Pope, "Cosmological Solutions, p-branes and the Wheeler De Witt Equation", [hep-th/9707182](#).
- [56] V.R. Gavrillov and V.N. Melnikov, Toda lattices with type  $A_n$  Lie algebras for multidimensional cosmology with intersecting  $p$ -branes. Proceedings of International Seminar on Multidimensional Cosmology (Potsdam, March 1998).